

The footprint of extra dimensions

- 4d supergravities appear when compactifying string theory
- Fluctuations of the internal geometry translates into **massless** 4d **scalar fields** known as “*moduli*”

$$\mathcal{L} = \frac{1}{2} R - \frac{1}{2} \partial_\mu \phi_i \partial^\mu \phi^i$$



Deviations
from GR !!

massless scalars = long range interactions (precision tests of GR)

Linking strings to observations \rightarrow Mechanisms to stabilise moduli !!


$$V(\phi) = m_{ij}^2 \phi^i \phi^j + \dots$$

‣ Moduli VEVs $\langle \phi \rangle = \phi_0$ determine 4d physics

$\left\{ \begin{array}{l} \Lambda_{c.c} \equiv V(\phi_0) \\ g_s \text{ and } \text{Vol}_{int} \\ \text{fermi masses} \end{array} \right.$

How to deform massless theories to have $V(\phi) \neq 0$?

- Supersymmetry dictates what deformations are allowed



gaugings = part of the global symmetry is promoted to local (*gauge*)

- Gauged supergravities can be systematically studied

embedding tensor formalism

[Nicolai, Samtleben '00]

[de Wit, Samtleben, Trigiante '02 '05 '07]


[Schon, Weidner '06]

undeformed = abelian = massless

- › Reducing 10d supergravities down to 4d

global symmetry G = **stringy features** (*i.e.* U/T/S-duality)

- › The **physical scalar sector** parameterises the **coset space** $\mathcal{M} = G/H$ where H is the maximal compact subgroup of G



- › **Abelian** gauge fields $A_{\mu}^{\mathcal{F}}$ in the fundamental of G

deformed = non-abelian = massive

- › *Gauging* : a non-abelian subgroup $G_0 \subset G$ is promoted to a **gauge symmetry** yielding a *gauged* supergravity

$$\nabla_\mu \longrightarrow \nabla_\mu - g A_\mu^{\mathcal{F}} \Theta_{\mathcal{F}}^{\mathcal{A}} t_{\mathcal{A}} \quad \text{where} \quad \begin{cases} \mathcal{F} \equiv \text{fund} \\ \mathcal{A} \equiv \text{adj} \end{cases} \text{ reps of } G$$

embedding tensor

Consistent gauge group \implies Quadratic constraints on Θ


- › **Non-trivial scalar potential** for the scalar fields $\mathcal{M} = G/H$

$$V(\Theta, \mathcal{M}) \neq 0$$

duality invariant = stringy !!

- › The embedding tensor $\Theta_{\mathcal{F}}^{\mathcal{A}}$ encodes **any possible deformation** of the massless theory **irrespective of its higher-dimensional origin**

A one minute summary



A chain of truncations in 4d

[Dibitetto, A.G, Roest in progress]

$G = E_7$
e.t comp = 912
vectors = 56
scalars = 133
 $\mathcal{N} = 8$

$SO(3)$ →

$G = SL(2) \times G_2$
e.t comp = 72
vectors = 4
scalars = 17
 $\mathcal{N} = 2$

↓ \mathbb{Z}_2

$G = SL(2) \times SO(6, 6)$
e.t comp = 464
vectors = 24
scalars = 69
 $\mathcal{N} = 4$

$SO(3)$ →

$G = SL(2) \times SL(2) \times SL(2)$
e.t comp = 40
vectors = 0
scalars = 9
 $\mathcal{N} = 1$

\mathbb{Z}_2 ↓

Half-maximal supergravities in 4d

[Schon, Weidner '06]

$$G = SL(2) \times SO(6, 6)$$

$$\text{e.t comp} = 464$$

$$\text{vectors} = 24$$

$$\text{scalars} = 69$$

$$\mathcal{N} = 4$$

Questions :

- › Can the whole vacuum structure be charted in $\mathcal{N} = 4$ theories ?
- › Are there connections in the landscape of vacua ?

Half-maximal : symmetry and fields

‣ Global symmetry (duality) group

$$G = SL(2) \times SO(6, 6)$$

‣ **Field content** = supergravity multiplet + six vector multiplets

‣ **Vectors** $A_{\mu}^{\alpha M}$ in the fundamental of G

$\alpha = +, -$ is an electric-magnetic $SL(2)$ index

$M = 1, \dots, 12$ is an $SO(6, 6)$ index

} **24 vectors**

‣ The **physical scalars** parameterise $\mathcal{M} = M_{\alpha\beta} \times M_{MN}$

• 1 axion + 1 dilaton in $SL(2)$

• 30 axions + 6 dilatons in $SO(6,6)$

} **38 physical scalars**

Half-maximal : *gaugings* and scalar potential

- *Gaugings* are classified by two **embedding tensor pieces**

$$\xi_{\alpha M} \in (\mathbf{2}, \mathbf{12}) \quad \text{and} \quad f_{\alpha MNP} \in (\mathbf{2}, \mathbf{220})$$

* reps of $SL(2) \times SO(6,6)$

- Supersymmetry + gauge invariance determine the **scalar potential**

$$V = \frac{1}{64} f_{\alpha MNP} f_{\beta QRS} M^{\alpha\beta} \left[\frac{1}{3} M^{MQ} M^{NR} M^{PS} + \left(\frac{2}{3} \eta^{MQ} - M^{MQ} \right) \eta^{NR} \eta^{PS} \right] -$$

$$- \frac{1}{144} f_{\alpha MNP} f_{\beta QRS} \epsilon^{\alpha\beta} M^{MNPQRS} + \frac{3}{64} \xi_{\alpha}^M \xi_{\beta}^N M^{\alpha\beta} M_{MN}$$

$\eta \equiv SO(6,6)$ -metric

To keep in mind : V is **quadratic** in the emb. tens. parameters

- 464 e.t. components + 38 physical scalars = **TOO MUCH !!**

The $SO(3)$ truncation

- Keeping only fields and embedding tensor components invariant under the action of a subgroup $SO(3) \subset SO(6,6)$

$$G = SL(2) \times SO(6,6)$$

$$\text{e.t comp} = 464$$

$$\text{vectors} = 24$$

$$\text{physical scalars} = 38$$

$$\mathcal{N} = 4$$

$$\xrightarrow{SO(3)}$$

$$G = SL(2) \times SL(2) \times SL(2)$$

$$\text{e.t comp} = 40$$

$$\text{vectors} = 0$$

$$\text{physical scalars} = 6$$

$$\mathcal{N} = 1$$

- R-symmetry group :

$$SU(4) \supset SO(3)$$

$$4 \longrightarrow 1 + 3$$

The $SO(3)$ truncation : fields and *gaugings*

[Derendinger, Kounnas, Petropoulos, Zwirner '04]

› Symmetry and fields :

- global symmetry $G = SL(2)_S \times SL(2)_T \times SL(2)_U$
- $A_\mu^{\alpha M} = 0$ and $\xi_{\alpha M} = 0$
- scalar coset = **3 complex scalars** = STU - models !!

$$\underbrace{S \equiv \chi + i e^{-\phi}}_{SL(2)_S}, \quad \underbrace{T \equiv \chi_1 + i e^{-\varphi_1} \quad \text{and} \quad U \equiv \chi_2 + i e^{-\varphi_2}}_{SL(2)_T \times SL(2)_U}$$

› The *gaugings* $G_0 \subset G$ and the scalar potential $V(S, T, U)$ are specified by the embedding tensor

$$f_{\alpha MNP} = 40 \text{ components}$$

Gaugings from fluxes

- › String embedding as flux compactification

$$f_{\alpha MNP} = \text{generalised fluxes}$$

Example: SO(3) truncation \longleftrightarrow type II orientifolds of $\mathbb{T}^6 / \mathbb{Z}_2 \times \mathbb{Z}_2$

- › Type IIB fluxes and embedding tensor

$$\begin{aligned} f_{+mnp} &= \tilde{F}'_{mnp} & , & & f_{+mn}{}^p &= Q'_{mn}{}^p & , & & f_{+}{}^{mn}{}_p &= Q^{mn}{}_p & , & & f_{+}{}^{mnp} &= \tilde{F}^{mnp} & , \\ f_{-mnp} &= \tilde{H}'_{mnp} & , & & f_{-mn}{}^p &= P'_{mn}{}^p & , & & f_{-}{}^{mn}{}_p &= P^{mn}{}_p & , & & f_{-}{}^{mnp} &= \tilde{H}^{mnp} & , \end{aligned}$$

[Dibitetto, Linares, Roest '10]

* index splitting $M = (m, {}^m)$

- › Perfect matching with flux-induced superpotential (**up to Q.C**)

$$W_{flux} = (P_F - P_H S) + 3T (P_Q - P_P S) + 3T^2 (P_{Q'} - P_{P'} S) + T^3 (P_{F'} - P_{H'} S)$$

Gaugings and sources

› Consistency of the gauge group = quadratic constraints on $f_{\alpha MNP}$

gaugings

$$A_\mu = A_\mu^{\alpha M} T_{\alpha M}$$

$$[T_{\alpha M}, T_{\beta N}] = f_{\alpha MN}{}^P T_{\beta P}$$



Quadratic Constraints

$$\epsilon^{\alpha\beta} f_{\alpha MNR} f_{\beta PQ}{}^R = 0$$

$$f_{\alpha R[MN} f_{\beta PQ]}{}^R = 0$$

› String theory :

quadratic constraints = Absence of sources breaking $\mathcal{N} = 4$

Example : Type IIB orientifolds with O3/O7-planes

- (H, F) fluxes : Unconstrained D3-brane flux-induced tadpole
- (H, F, Q) fluxes : Vanishing of D7-brane flux-induced tadpole
- (H, F, Q, P) fluxes : Vanishing of D7, NS7 and I7 flux-induced tadpoles
- ??

We would like to . . .

- 1) build **all** the consistent SO(3)-invariant gaugings specified by $f_{\alpha MNP}$ by solving the quadratic constraints

$$\epsilon f f = 0 \quad \text{and} \quad f f = 0$$

- 2) compute **all** the SO(3)-invariant extrema of the f -induced scalar potential $V(f, \Phi)$ by solving the extremisation conditions


$$\left. \frac{\partial V}{\partial \Phi} \right|_{\Phi_0} = 0 \quad \text{with} \quad \Phi \equiv (S, T, U)$$

- 3) check stability of these extrema with respect to fluctuations of **all** the 38 scalars of half-maximal supergravity
- 4) identify the gauge group G_0 underlying **all** the different solutions

. . . but is this doable ?

Strategy and tools

- **Idea** : use the global symmetry group (non-compact part) to bring **any** field solution back to the origin !!



- At the origin everything is **simply quadratic** in the $f_{\alpha MNP}$ parameters

$$V(\Phi) = \sum_{\text{terms}} f f \Phi^{\text{high degree}}$$


so then,

computing the
vacua structure

=

multivariate polynomial system

$$I = \langle \partial_{\Phi} V|_{\Phi_0}, \epsilon f f, f f \rangle$$



Algebraic Geometry techniques !!

Basics of Algebraic Geometry



- Algebraic Geometry studies multivariate polynomial system and their link to geometry

$$I = \langle P_1, P_2 \rangle$$

$$P_1(x, y, z) = xz$$

$$P_2(x, y, z) = yz$$


algebraic system



variety

- GTZ prime decomposition (analogous to integers dec. $30 = 2 \times 3 \times 5$)

$$I = J_1 \cap J_2 \quad \text{where} \quad \begin{cases} J_1 = \langle z \rangle \longrightarrow xy\text{-plane} \\ J_2 = \langle x, y \rangle \longrightarrow z\text{-axis} \end{cases}$$



$$J_1 \cap J_2 \longleftrightarrow V(J_1) \cup V(J_2)$$

algebra-geometry dictionary

- Applying the above procedure to our problem involving fluxes

$$I = \langle \partial_\Phi V|_{\Phi_0}, \epsilon f f, f f \rangle$$

$$I = J_1 \cap J_2 \cap \dots \cap J_n$$



Splitting of the landscape
into n disconnected pieces !!

An example : type IIA with metric fluxes

- › Testing the method with type IIA orientifold models including **gauge fluxes** and a **metric flux**


[Dall'Agatta, Villadoro, Zwirner '09]

$$\left(F_{p=0,2,4,6} , H_3 \right) + \omega \subset f_{\alpha MNP}$$

- › **Subset** of embedding tensor components **closed under** $G_{n,c}$
 - ✓ Fields can still be set at the origin without lost of generality
 - ✓ Stability with respect to fluctuations around the origin can be computed

[Borghese, Roest '10]

- › **Vacua structure** of these type IIA orientifolds



[Giddings, Kachru, Polchinski '02]


➤ An AdS_4 landscape

$$16 = 4 + 4 + 4 + 4$$

➤ Unique gauging

$$G_0 = ISO(3) \times U(1)^6$$

$1_{(\pm,\pm)}$	$2_{(\pm,\pm)}$	$3_{(\pm,\pm)}$	$4_{(\pm,\pm)}$
$\mathcal{N} = 1$ SUSY & SUSY	SUSY	SUSY	SUSY
stable	unstable	stable	stable
$m^2 = -2/3$	$m^2 = -4/5$	$m^2 > 0$	$m^2 > 0$
$V = -1$	$V = -32/27$	$V = -8/15$	$V = -32/27$



(* $m^2 \equiv$ lightest mode (B.F. bound = $-3/4$)

➤ All the solutions are connected and correspond to $\omega \equiv SU(2) \times SU(2)$

[Caviezel, Koerber, Körs, Lüst, Tsimpis/Wrase, Zagermann '08, '08]

NO SOURCES AT ALL !!

Lifting to maximal supergravity ?

$$G = E_7$$

e.t comp = 912
vectors = 56
scalars = 133

$$\mathcal{N} = 8$$

↑ \mathbb{Z}_2

$$G = SL(2) \times SO(6, 6)$$

e.t comp = (2,12) + (2,220)
vectors = (2,12)
scalars = (3,1) + (1,66)

$$\mathcal{N} = 4$$

* reps of G

Question :

- Is the absence of sources enough for the geometric IIA solutions to lift to a maximal supergravity theory?

Half-maximal inside maximal

[Dibitetto, A.G., Roest '11]

- Look at the maximal theory with the half-maximal ``glasses''

$$\begin{array}{l}
 E_7 \quad \supset \quad SL(2) \times SO(6,6) \\
 \text{vectors : } 56 \longrightarrow (2, 12) + (1, 32) \\
 \text{scalars : } 133 \longrightarrow (3, 1) + (1, 66) + (2, 32') \\
 \text{e.t: } \underbrace{912}_{X_{56} \ 56} \longrightarrow \underbrace{(2, 12)}_{\xi_{\alpha M}} + \underbrace{(2, 220)}_{f_{\alpha MNP}} + (1, 352') + (3, 32)
 \end{array}
 \left. \vphantom{\begin{array}{l} \\ \\ \\ \end{array}} \right\}
 \begin{array}{l}
 \mathbb{Z}_2 \\
 \text{rep} \equiv \text{bos} (B) \rightarrow \text{even} \\
 \text{rep} \equiv \underbrace{\text{fermi} (F)}_{SO(6,6)} \rightarrow \text{odd}
 \end{array}$$

- Gauge algebra in the maximal theory

$$\left. \begin{array}{l}
 [A_B, A_B] = X_{BB}^B A_B + X_{BB}^F A_F \\
 [A_B, A_F] = X_{BF}^B A_B + X_{BF}^F A_F \\
 [A_F, A_F] = X_{FF}^B A_B + X_{FF}^F A_F
 \end{array} \right\} \text{Jacobi identities} = QC_{\mathcal{N}=8}$$

- Components with an **odd** number of **fermionic indices** projected out !!

Extra conditions for the lifting

- For a half-maximal to be embeddable in a maximal theory

$$QC_{\mathcal{N}=8} = QC_{\mathcal{N}=4} + \text{extra conditions for the lifting}$$

- The **extra conditions** are

$$\underbrace{f_{\alpha MNP} f_{\beta}{}^{MNP} = 0}_{(3,1)} \quad \text{and} \quad \underbrace{\epsilon^{\alpha\beta} f_{\alpha[MNP} f_{\beta QRS]} \Big|_{\text{SD}} = 0}_{(1,462')}$$

* reps of $SL(2) \times SO(6,6)$

Example: type IIB dual sources ?

$$\underbrace{H_3 \wedge F_3}_{\text{D3-brane flux-induced tadpole}} \subset (1,462') \quad \longrightarrow \quad \text{which objects fill the rep ?}$$

An example : lifting of geometric type IIA

- **All** the 16 geom. IIA solutions **lift** to maximal supergravity

- Fake SUSY becomes SUSY

$$\underbrace{SU(8)}_{\mathcal{N}=8 \text{ SUSY}} \rightarrow \underbrace{SU(4)}_{\mathcal{N}=4 \text{ SUSY}} \times \underbrace{SU(4)}_{\mathcal{N}=4 \text{ FAKE}}$$

$1_{(\pm,\pm)}$	$2_{(\pm,\pm)}$	$3_{(\pm,\pm)}$	$4_{(\pm,\pm)}$
$\mathcal{N} = 1 \text{ SUSY}$	SUSY	SUSY	SUSY
stable	unstable	stable	unstable
$m^2 = -2/3$	$m^2 = -4/5$	$m^2 > 0$	$m^2 = -4/3$
$V = -1$	$V = -32/27$	$V = -8/15$	$V = -32/27$

(* $m^2 \equiv$ lightest mode (B.F. bound = $-3/4$)

- Masses of the 70 physical scalars \Rightarrow ~~SUSY~~ and **stable** minimum !!

[Le Diffon, Samtleben, Trigiante '11]

- Underlying gaugings in maximal supergravity (28 vectors) ??

(...in progress)

Conclusions

- Some progress towards disentangling the landscape of extended supergravities can still be done without performing statistics of vacua
- The approach relies on the combined use of global symmetries and of algebraic geometry techniques
- As a warming-up, the complete vacua structure of simple type IIA orientifold theories can be worked out revealing some odd features :
 - i)* connections between vacua
 - ii)* $\mathcal{N} = 8$ lifting of the entire vacua structure
 - iii)* stability without supersymmetry
- **For the future :**
 - Beyond the geometric limit : non-geometric backgrounds, dual branes . . .
 - de Sitter in extended supergravity (maybe $\mathcal{N} = 2$) and links to Cosmology
 - Higher dimensional origin of gaugings : DFT, Generalised Geometry . . .



Thanks !!

Higher-dimensional origin of gaugings

guiding principle = duality covariance

Example: T-duality = SO(d,d)

➤ **DFT / Doubled Geometry** $\Rightarrow \{ \partial_m , \partial^m \}$
(fundamental = 1-form)

[Hull, Zwiebach '09]
[Hohm, Hull, Zwiebach '10]
[Hohm, Kwak, Zwiebach '11]
[Geissbühler '11]
[Aldazabal, Baron, Marques, Nunez '11]

➤ **Generalised Geometry** $\Rightarrow \{ G_{mn} , B_{mn} , \beta^{mn} \}$
(adjoint = 2-form)

[Gualtieri '04]
[Grana, Minasian, Petrini, Tomasiello '04 '05]
[Grana, Minasian, Petrini, Waldram '08]
[Coimbra, Strickland-Constable, Waldram '11]

➤ **Non-Geometry** $\Rightarrow \{ H_{mnp} , \omega_{np}^m , Q_p^{mn} , R^{mnp} \}$
(3-form)

[Shelton, Taylor, Wecht '05 '06]
[Aldazabal, Cámara, Font, Ibañez '06]
[de Carlos, A.G. Moreno '09]
[Aldazabal, Andrés, Cámara, Grana '10]