

Advanced General Relativity : Exercises sheet (2020-2021)

Exercise 1

In a space-time with torsion, one has that

$$\Gamma_{\mu\nu}{}^\rho = \Gamma_{\mu\nu}{}^\rho(g) - K_{\mu\nu}{}^\rho, \quad (1)$$

where $\Gamma_{\mu\nu}{}^\rho(g)$ are the torsion-free Christoffel symbols and $K_{\mu\nu}{}^\rho$ is the so-called *contorsion* tensor. Show that

$$\int d^4x \sqrt{-|g|} \nabla_\mu V^\mu, \quad (2)$$

is *not* a total derivative. Instead, it can be expressed as a boundary term plus an extra piece depending on the contorsion tensor, thus spoiling the usual integration by parts.

Note: $\partial_\mu \sqrt{-|g|} = \sqrt{-|g|} \Gamma_{\rho\mu}{}^\rho(g)$

Exercise 2

Starting from the action for a spin 1 (Maxwell) field

$$S_A = \int d^4x \sqrt{-|g|} \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right), \quad (3)$$

with field strength

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu, \quad (4)$$

and assuming a space-time without torsion :

- Compute the equation of motion for the Maxwell field A_μ .
- Compute the energy momentum tensor $T_{\mu\nu}$.
- Compute $T \equiv T_\mu{}^\mu$ and $\nabla_\mu T^{\mu\nu}$.

Exercise 3

Let us consider a simple wormhole metric of the form

$$ds^2 = -dt^2 + du^2 + (b_0^2 + u^2)(d\theta^2 + \sin^2\theta d\phi^2), \quad (5)$$

in terms of a constant parameter $b_0 > 0$. The ranges of the coordinates are given by

$$t \in (-\infty, \infty) \quad , \quad u \in (-\infty, \infty) \quad , \quad \theta \in [0, \pi] \quad , \quad \phi \in [0, 2\pi] . \quad (6)$$

- Using cylindrical coordinates in the embedding space

$$ds^2 = dz^2 + dr^2 + r^2 d\phi^2, \quad (7)$$

construct an embedding diagram

$$z(r) \quad \text{with} \quad r^2 = b_0^2 + u^2, \quad (8)$$

for the equatorial plane $\theta = \pi/2$ at a fixed time t . How many asymptotic regions does the geometry have?

- Using the Einstein equations, compute the stress tensor $T_{\mu\nu}$ compatible with the metric (5). What kind of matter would this stress tensor be accounting for? Is it an ordinary type of matter?

Exercise 4

Consider a field theory including a spin 2 field (metric) and a massive spin 3/2 field (gravitino) in the presence of a cosmological constant. The action of the theory is given by

$$S = S_g + S_\Lambda + S_\Psi + S_{\text{mass}} , \quad (9)$$

with

$$\begin{aligned} S_g &= \frac{1}{2\kappa^2} \int d^4x e e_a{}^\mu e_b{}^\nu R_{\mu\nu}{}^{ab} , \\ S_\Lambda &= -\frac{1}{2\kappa^2} \int d^4x e \Lambda , \\ S_\Psi &= -\frac{1}{2\kappa^2} \int d^4x e \bar{\Psi}_\mu \gamma^{\mu\nu\rho} D_\nu \Psi_\rho , \\ S_{\text{mass}} &= -\frac{1}{2\kappa^2} \int d^4x e m \bar{\Psi}_\mu \gamma^{\mu\nu} \Psi_\nu . \end{aligned} \quad (10)$$

The first contribution S_g is the usual Einstein-Hilbert action for the metric. The second contribution S_Λ is that of a cosmological constant Λ . The third contribution S_Ψ is the Rarita-Schwinger action for the gravitino field. Finally, the fourth contribution S_{mass} with constant m is just a mass term for the gravitino field.

Show that the above action S is invariant, to lowest order in fermions, under the local supersymmetry transformations

$$\begin{aligned} \delta_\epsilon e_\mu{}^a &= \frac{1}{2} \bar{\epsilon} \gamma^a \Psi_\mu , \\ \delta_\epsilon \Psi_\mu &= D_\mu \epsilon - g \gamma_\mu \epsilon , \end{aligned} \quad (11)$$

with (spinorial) supersymmetry parameter $\epsilon_\alpha(x)$ provided two relations of the form

$$\Lambda = c_1 c_2 g^2 \quad , \quad m = c_2 g , \quad (12)$$

hold with c_1 and c_2 being constants. More concretely:

- a) Determine the values of c_1 and c_2 .
- b) Discuss the relation between supersymmetry and the sign of the cosmological constant.

Note: $\gamma^{\mu\nu\rho} \gamma_\rho = 2 \gamma^{\mu\nu}$.

Note: $\bar{\chi} \gamma_{\mu_1 \mu_2 \dots \mu_n} \lambda = t_n \bar{\lambda} \gamma_{\mu_1 \mu_2 \dots \mu_n} \chi$ with $t_0 = t_3 = -t_1 = -t_2 = 1$.

Exercise 5

Let us consider two real scalar fields ϕ^1 and ϕ^2 serving as coordinates on a two-dimensional field space with metric

$$ds^2 = K_{ij}(\phi) d\phi^i d\phi^j = \frac{1}{(\phi^2)^2} \left((d\phi^1)^2 + (d\phi^2)^2 \right) . \quad (13)$$

Assuming a flat Minkowski space-time $g_{\mu\nu} = \eta_{\mu\nu}$ and taking the scalar fields to be only a function of time, *i.e.* $\phi^i = \phi^i(t)$:

a) Show that the action for the scalar fields

$$S = \int d^4x \left(-\frac{1}{2} K_{ij} \partial_\mu \phi^i \partial^\mu \phi^j \right), \quad (14)$$

takes the simple form

$$S = \frac{1}{2} \int dt \frac{1}{(\phi^2)^2} \left((\dot{\phi}^1)^2 + (\dot{\phi}^2)^2 \right), \quad (15)$$

where we have denoted $\dot{\phi}^i = \frac{d\phi^i}{dt}$.

b) Show that the Euler-Lagrange equations of motion are given by

$$\ddot{\phi}^1 - \frac{2}{\phi^2} \dot{\phi}^1 \dot{\phi}^2 = 0 \quad \text{and} \quad \ddot{\phi}^2 + \frac{1}{\phi^2} \left((\dot{\phi}^1)^2 - (\dot{\phi}^2)^2 \right) = 0. \quad (16)$$

c) Show that the Euler-Lagrange equations (16) can be expressed as a geodesic equation in field space

$$\ddot{\phi}^i + \Gamma_{jk}{}^i \dot{\phi}^j \dot{\phi}^k = 0, \quad (17)$$

in terms of the Christoffel symbols in field space

$$\Gamma_{jk}{}^i = \frac{1}{2} K^{il} (\partial_j K_{lk} + \partial_k K_{jl} - \partial_l K_{jk}). \quad (18)$$

d) Can you identify the coset space $\text{SL}(2)/\text{SO}(2)$ in (13) upon a suitable field redefinition (change of coordinates in field space)?

Exercise 6

Let us consider the group of diffeomorphisms in a $(D+1)$ -dimensional theory of gravity with space-time coordinates $x^M = (x^\mu, z)$. These diffeomorphisms are given by transformations on the metric of the form

$$\delta_{\widehat{\xi}} \widehat{g}_{MN} = \widehat{\xi}^P \partial_P \widehat{g}_{MN} + \widehat{g}_{MP} \partial_N \widehat{\xi}^P + \widehat{g}_{PN} \partial_M \widehat{\xi}^P \quad (19)$$

where

$$\widehat{\xi}^M(x, z) = \left(\widehat{\xi}^\mu, \widehat{\xi}^z \right), \quad (20)$$

is the $(D+1)$ -dimensional diffeomorphism parameter and

$$\widehat{\xi}^\mu = \xi^\mu(x), \quad \widehat{\xi}^z(x, z) = \theta(x) + cz, \quad (21)$$

with $\theta(x)$ being an arbitrary function of x^μ and c being a constant. In addition to the above diffeomorphisms, let us also consider scaling transformations on the metric of the form

$$\delta_a \widehat{g}_{MN} = 2a \widehat{g}_{MN}, \quad (22)$$

and a being a real constant.

a) Using the Kaluza-Klein Ansatz for the $(D+1)$ -dimensional metric

$$\widehat{g}_{MN} = \begin{pmatrix} e^{2\alpha\phi} g_{\mu\nu} + e^{2\beta\phi} A_\mu A_\nu & e^{2\beta\phi} A_\mu \\ e^{2\beta\phi} A_\mu & e^{2\beta\phi} \end{pmatrix}, \quad (23)$$

with $\beta = -(D-2)\alpha$, show that (19) and (22) induce transformations $\delta = \delta_{\widehat{\xi}} + \delta_a$ on the D -dimensional fields of the form

$$\begin{aligned}\delta\phi &= \xi^\rho \partial_\rho \phi - \frac{1}{(D-2)\alpha} (c+a), \\ \delta A_\mu &= \xi^\rho \partial_\rho A_\mu + A_\rho \partial_\mu \xi^\rho + \partial_\mu \theta - c A_\mu, \\ \delta g_{\mu\nu} &= \xi^\rho \partial_\rho g_{\mu\nu} + g_{\mu\rho} \partial_\nu \xi^\rho + g_{\rho\nu} \partial_\mu \xi^\rho + \frac{2}{(D-2)} \left[c + a(D-1) \right] g_{\mu\nu}.\end{aligned}\tag{24}$$

b) Discuss the transformations arising upon the particular choices of parameters $a = -\frac{c}{(D-1)}$ and $a = -c$.

Exercise 7

Let us consider a Maxwell field \widehat{B}_M in $D+1$ dimensions, with a $(D+1)$ -dimensional index $M = \mu \oplus z$ splitting into a D -dimensional index μ and an additional direction z . The $(D+1)$ -dimensional Maxwell field \widehat{B}_M can then be decomposed as

$$\widehat{B}_M = (\widehat{B}_\mu, \widehat{B}_z) \equiv (B_\mu(x), \chi(x)).\tag{25}$$

Using the Kaluza-Klein Ansatz for the $(D+1)$ -dimensional frame and its inverse

$$\widehat{e}_M^A = \begin{pmatrix} e^{\alpha\phi} e_\mu^a & e^{\beta\phi} A_\mu \\ 0_{1\times 4} & e^{\beta\phi} \end{pmatrix}, \quad \widehat{e}_A^N = \begin{pmatrix} e^{-\alpha\phi} e_a^\nu & -e^{-\alpha\phi} A_a \\ 0_{1\times 4} & e^{-\beta\phi} \end{pmatrix},\tag{26}$$

where $A_a \equiv e_a^\nu A_\nu$:

- Show that $\widehat{e}_M^A \widehat{e}_A^N = \delta_M^N$.
- Compute $\widehat{g}_{MN} = \widehat{e}_M^A \widehat{e}_N^B \widehat{\eta}_{AB}$, where $\widehat{\eta}_{AB}$ is the $(D+1)$ -dimensional Minkowski metric, and show that the result agrees with (23).
- Show that, when $\beta = -(D-2)\alpha$, the $(D+1)$ -dimensional Maxwell action

$$S_{\widehat{B}} = \int d^{D+1}x \sqrt{-|\widehat{g}|} \left(-\frac{1}{4} \widehat{F}_{MN} \widehat{F}^{MN} \right) = \int d^{D+1}x \widehat{e} \left(-\frac{1}{4} \widehat{F}_{AB} \widehat{F}^{AB} \right),\tag{27}$$

reduces to a D -dimensional Maxwell-scalar action of the form

$$\begin{aligned}S_{\widehat{B}} &= (2\pi L) \int d^Dx e \left(-\frac{1}{4} e^{-2\alpha\phi} \mathcal{F}_{ab} \mathcal{F}^{ab} - \frac{1}{2} e^{2(D-2)\alpha\phi} \partial_a \chi \partial^a \chi \right), \\ &= (2\pi L) \int d^Dx \sqrt{-|g|} \left(-\frac{1}{4} e^{-2\alpha\phi} \mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu} - \frac{1}{2} e^{2(D-2)\alpha\phi} \partial_\mu \chi \partial^\mu \chi \right),\end{aligned}\tag{28}$$

with L being the radius of the (circle) z -direction, and where we have defined

$$\partial_a \equiv e_a^\nu \partial_\nu, \quad \mathcal{F}_{ab} \equiv F_{ab} - 2 \partial_{[a} \chi A_{b]}.\tag{29}$$